## Summary of Phase I WIP Loads - Worcester

The charts in this handout summarize the Phase I WIP loads by major source sector. Although these numbers are expected to be updated in early summer 2011 by the US EPA, they are being provided in response to general interest by the Phase II WIP Local Teams. The following footnotes are important. There is a near certainty that these numbers will change. Therefore strategies developed on the basis of these numbers may need to be revised when final updated allocations are received.

1. Loads are reported as delivered loads to the Bay. Note that delivery factors (i.e. the proportion of load delivered to the Bay) can change between scenarios.

2. EPA's revised Bay watershed model will significantly revise how agricultural nutrient management plans are addressed, and will increase the amount of low density urban.

- 3. Scenarios presented<sup>1</sup>:
  - 2009 = Progress
  - 2017 Strategy = Result of Phase I WIP strategies
  - 2017 Target = 70% from 2009 Progress to 2020 Target for nonpoint sources (including stormwater)
  - 2020 Target = Allocation of the TMDL
  - -- The Final point source loads are the cap strategy and not the estimated load in 2020. The estimated load in 2020 is presented as "2020 Projected".

4. The State reserves the right to adjust the remaining WWTP load allocation and/or make other necessary adjustments if the state is falling short of its overall goals.

5. The 2017 allocated loads and target reductions assume full implementation of ENR and other upgrades by the WWTP sector. These upgrades accomplish a disproportionately large share of the 70% statewide target reduction set for 2017, compared to other sectors.

6. The 2017 Target is to achieve relatively evenly distributed progress in all source sectors toward their final nonpoint source targets between now and 2020. The 2020 targets, then, should be the focus of local planning and implementation strategies, including 2-year milestones.

7. CAFO (Concentrated Animal Feeding Operation) loads are associated with the animal production area, which is why the loads are small. Manure that is applied on fields is counted within the Agricultural sources.

8. Non-Regulated (Non-MS4) Stormwater (SW) reductions are associated with additional fertilizer management (urban nutrient management) on commercially managed lawns.

9. "Urban Regulated" includes all Urban load for Phase I and Phase II NPDES Permitted jurisdictions and for Construction, Extractive, and Industrial lands.





**87,875 - 100,887 115.170 - 138.446 — 140,189 — 154,91**0 194.590 - 220.688

TP (lb/yr):

**14.845 - 77.412** 

## Worcester WIP 2017 Strategies

#### **Point Sources:**

• Major WWTPs – Continue Upgrades to ENR

• Minor Industrial - Identify loading targets and issue schedules in permits for reductions

 Sewer Overflows were eliminated in 2006 - the CSO load shown in the charts on the following pages should be zero

#### Urban:

 Urban Nutrient Management Law – Continue the regulation of fertilizer applications on commercially managed lawns

### Septics:

• Continue Upgrade of new and failing Septic Systems in the Critical Area

• Septic hookups to ENR plants

• Require upgrade of all systems in Critical Area

### Agriculture:

• Various Practices (i.e. Cover Crops, Conservation Plans, Nutrient Management Plans, Forest/Grass Buffers, and Animal Waste Management BMPs on AFO (Animal Feeding Operations) and CAFO (Concentrated Animal Feeding Operations))

Air:

- Clean Air Act Implementation
- Maryland Healthy Air Act
- Clean Air Interstate Rule
- Tier-2 Vehicle Rule
- Nonroad Engine Rule
- Heavy-Duty Diesel Engine Rule
- Locomotive/Marine Engine Rule





\* Different assumptions were used by the US EPA for 2009 septic loads versus 2017 and 2020. Reductions from 2009 are likely less than indicated here.



Printed on: 4/5/2011



# Worcester Total Nitrogen Loads (lb/yr - Delivered) - A Closer Look at Source Sector Categories



# Worcester Total Phosphorus Loads (lb/yr - Delivered) - A Closer Look at Source Sector Categories